TABLE 4 (continued) MONITORING PROGRESS TOWARD ARIZONA AND SELECTED NATIONAL YEAR 2010 OBJECTIVES

Focus areas and selected objectives: (in parentheses are <i>Healthy People 2010</i> objective numbers)	Baseline ARIZONA RATES OR CASES BY YEAR:												
	for the U.S.	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2010 TARGET
4D. INJURY AND VIOLENCE													
Reduce firearm-related deaths (HP15-3)	11.3	15.3	15.5	17.3	14.8			>		>			4.1
Reduce deaths caused by poisonings (HP15-8)	6.8	7.6	11.7	12.6	12.0	>		>	•	>	•	>	1.5
Reduce deaths caused by suffocation (HP15-9)	4.1	4.2	4.5	4.5	5.0			>			—		3.0
Reduce deaths caused by unintentional injuries (HP15-13)	35.0	41.1	46.6	45.2	44.8	>	•	•	•	•	•	•	17.5
Reduce deaths caused by motor vehicle crashes (HP15-15)	15.6	17.5	17.2	18.5	18.0	>	•	•	•	•	•	•	♦ 16.0
Reduce residential fire deaths (HP15-25)	1.2	0.9	0.9	0.8	0.7	>	•					>	0.2
Reduce deaths from falls (HP15-27)	4.7	7.6	8.9	8.6	8.8	>		•			>	>	3.0
♦ Reduce drownings (HP15-29)	1.6	1.7	1.6	1.2	1.5	>	>	•	>			>	• 1.0
♦ Reduce homicides (HP15-32)	6.5	7.6	8.7	8.6	8.3	>			>				♦ 5.0
♦ Reduce the suicide rate (HP18-1)	11.3	14.6	11.5	15.9	14.6	•				—			♦ 10.0
Reduce the suicide rate among adolescents aged 15 to 19 years	9.4	11.1	10.5	9.9	9.7	>	>	•	>	•	>	>	♦ 10.0
Reduce the suicide rate among older adults aged 65 years and older	16.8	23.4	16.4	23.2	19.8	>	♦ 10.0						
4E. CANCER													
Reduce the overall cancer death rate (HP3-1)	202.4	170.4	169.2	168.6	169.4	•	•	•	•	•	•		159.9
Reduce the lung cancer death rate (HP3-2)	57.6	46.1	44.5	46.6	47.4	•	>	•	>		•	>	44.9
Reduce the breast cancer death rate (HP3-3)	27.9	25.4	23.6	22.2	22.1		22.3						
Reduce the death rate from cancer of the uterine cervix (HP3-4)	3.0	3.1	2.8	2.2	2.2	>	>	•	>	•	>	•	2.0
Reduce the colorectal cancer death rate (HP3-5)	21.2	17.0	16.7	16.5	17.9	•		>				>	13.9
Reduce the oropharyngeal cancer death rate (HP3-6)	3.0	2.1	2.2	2.2	2.0	>	•	•	•	•	•	•	2.7
Reduce the prostate cancer death rate (HP3-7)	32.0	28.4	24.3	24.8	25.1		>		•	•	•	>	28.8
Reduce the rate of melanoma cancer deaths (HP3-8)	2.8	2.8	2.4	2.9	2.7	•	•	•	•	•	•	•	2.5

• Indicates objectives and targets identified in *Healthy Arizona 2010*.

Note: Firearm-related deaths, deaths caused by poisonings, and deaths caused by suffocation include accidents, homicides, and suicides. Residential fire deaths, deaths from falls, and drowning deaths include only accidents. All mortality rates are per 100,000 persons. All mortality rates that are not age-specific, are age-adjusted to the 2000 U.S. standard.

Note: The baseline year for the U.S. is 1997.